

Catalog Number: IR106-P2

Anti-TNF alpha antibody

Package size: 25, 100 μ l


Store at: -20°C

MW (kDa): 26-34, 17


Overview

| | |
|---|---|
| Product Name | TNF alpha antibody |
| Product Number | IR106-P2 |
| Gene Description | tumor necrosis factor |
| Clonality | Polyclonal |
| Host | Rabbit |
| Species Reactivity | Human, Mouse |
| Recommended Applications Dilutions | Western Blot 1:500-1:1500 Immunofluorescence 1:200 – 1:300 Immunohistochemistry (Paraffin) 1:100 – 1:200 |
| Storage Buffer | 100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative |
| Concentration | 1.51 mg/ml |
| Purity | Affinity column purified |
| Storage | Store at +4°C for short term storage. Long time storage is recommended at -20°C |
| Notes | Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. |


IR106-P2 anti-TNF alpha antibody
WB image



IR106-P2 anti-TNF alpha antibody
IHC image



IR106-P2 anti-TNF alpha antibody
ICC/IF image



Anti- TNF alpha antibody at 1/700 dilution

Lysates at 50 μ g per lane. This blot was produced using a 12% SDS-PAGE. Nitrocellulose membrane was then blocked for an hour before being incubated with IR106-P2 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling TNF alpha with IR106-P2 at 1/150.

Cells were fixed with 4% PFA for 10 min at RT, permeabilized with 0.1% NP-40 for 10 min at RT then blocked with 5% BSA for 30 min at RT. Cells were stained with IR106-P2 anti- TNF alpha antibody (red) at 1:200 and 4°C. DAPI (blue) was used as the nuclear counter stain.

IReal Biotechnology Co., Ltd.