

Catalog Number: IR102-426


## Anti-Survivin / BIRC5 antibody

Package size: 25, 100  $\mu$ l  
Store at: -20°C  
MW (kDa): 17


### Overview

<b>Product Name</b>	Survivin / BIRC5 antibody
<b>Product Number</b>	IR102-426
<b>Gene Description</b>	baculoviral IAP repeat containing 5
<b>Clonality</b>	Polyclonal
<b>Host</b>	Rabbit
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Recommended Applications Dilutions</b>	Western Blot 1:2000-1:3000 Immunofluorescence 1:400 – 1:800 Immunohistochemistry (Paraffin) 1:100 – 1:300
<b>Storage Buffer</b>	100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative
<b>Concentration</b>	1.42 mg/ml
<b>Purity</b>	Affinity column purified
<b>Storage</b>	Store at +4°C for short term storage. Long time storage is recommended at -20°C
<b>Notes</b>	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.


IR102-426 anti-Survivin / BIRC5 antibody WB image



IR102-426 anti-Survivin / BIRC5 antibody ICC/IF image



IR102-426 anti-Survivin / BIRC5 antibody IHC image



**Survivin / BIRC5 antibody at 1/2000 dilution.**  
Lysates at 40  $\mu$ g per lane. This blot was produced using a 15% SDS-PAGE. Nitrocellulose membrane was then blocked for an hour before being incubated with IR102-426 overnight at 4°C.

Cells were fixed with 4% PFA for 10 min at RT, permeabilized with 0.1% NP-40 for 10 min at RT then blocked with 5% BSA for 30 min at RT. anti- Survivin / BIRC5 antibody (red) at 1:300 and 4°C. DAPI (blue) was used as the nuclear counter stain.

Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling Survivin / BIRC5 with IR102-426 at 1/150.

iReal Biotechnology Co., Ltd.