

Catalog Number: IR238-892


Anti-Cytokeratin 20 antibody

Package size: 25, 100 μ l
Store at: -20°C
MW (kDa): 43-55


Overview

| | |
|---|---|
| Product Name | Cytokeratin 20 antibody |
| Product Number | IR238-892 |
| Gene Description | keratin 20, type II |
| Clonality | Polyclonal |
| Host | Rabbit |
| Species Reactivity | Human |
| Recommended Applications Dilutions | Western Blot 1:1000 – 1:2000 Immunofluorescence 1:200 – 1:500 Immunohistochemistry (Paraffin) 1:200 – 1:300 |
| Storage Buffer | 100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative |
| Concentration | 1.33 mg/ml |
| Purity | Affinity column purified |
| Storage | Store at +4°C for short term storage. Long time storage is recommended at -20°C |
| Notes | Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. |

IR238-892 anti-Cytokeratin 20 antibody WB image



IR238-892 anti-Cytokeratin 20 antibody IHC image




Anti-Cytokeratin 20 antibody - at 1/2000 dilution

Lysates/proteins at 45 μ g per lane.
This blot was produced using a 10% SDS-PAGE. Nitrocellulose membrane was then blocked an hour before being incubated with IR238-892 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling Cytokeratin 20 with IR238-892 at 1/200.

Image

IR238-892 anti-Cytokeratin 20 antibody ICC/IF image



Immunofluorescence: cells were fixed with 4% paraformaldehyde for 10 min at RT, permeabilized with 0.1% NP-40 for 10 min at RT then blocked with 5% BSA for 30 min at room temperature. Cells were stained with IR238-892 anti-Cytokeratin 20 antibody (red) at 1:200 and 4°C. DAPI (blue) was used as the nuclear counter stain.

IReal Biotechnology Co., Ltd.

 **iReal Antibodies**



**ISO 13485:2016 Quality
Management System
Certified**

 www.irealbio.com

 03 - 5260005

 irealbio@irealbio.com