

Catalog Number: IR58-200


## Anti-FAK / PTK2 antibody

Package size: 25, 100  $\mu$ l  
Store at: -20°C  
MW (kDa): 125


### Overview

<b>Product Name</b>	<b>FAK / PTK2 antibody</b>
<b>Product Number</b>	<b>IR58-200</b>
<b>Gene Description</b>	protein tyrosine kinase 2
<b>Clonality</b>	Polyclonal
<b>Host</b>	Rabbit
<b>Species Reactivity</b>	Human, mouse, rat
<b>Recommended Applications Dilutions</b>	Western Blot 1:500 – 1:1000 Immunofluorescence 1:200 – 1:500 Immunohistochemistry (Paraffin) 1:200 – 1:300
<b>Storage Buffer</b>	100mM Tris Glycine, 1% BSA, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative
<b>Concentration</b>	0.54 mg/ml
<b>Purity</b>	Affinity column purified
<b>Storage</b>	Store at +4°C for short term storage. Long time storage is recommended at -20°C
<b>Notes</b>	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

IR58-200 anti-FAK / PTK2 antibody WB image



IR58-200 anti-FAK / PTK2 antibody IHC image



#### Anti-FAK / PTK2 antibody at 1/500 dilution

Lysates at 30  $\mu$ g per lane. This blot was produced using a 7.5% SDS-PAGE. Nitrocellulose membrane was then blocked with 3% BSA for an hour before being incubated with IR58-200 overnight at 4°C.

Immunohistochemical of Paraffin sections.  
Sample: mouse cervix.  
FAK / PTK2 antibody (IR58-200) : 1-200.